

of granular cell plasma membranes are also average values and comparisons with the plasma membranes of other types of cell whose lipid compositions are maintained in a "steady state" are not valid. The phospholipids and probably the triglycerides as well are completely catabolized as the differentiation of the epidermal cell is completed and it passes from the upper granular layer into the stratum corneum. There is also a partial loss of glycolipids [1]. We have suggested that our preparations of plasma membranes represent those from cells in the upper granular layer of the epidermis. On this basis the relative proportions of the "stable" or "retained" lipids in these membranes, that is, of cholesterol, fatty acids, ceramides, the remaining glycolipids and small amounts of cholesterol esters, hydrocarbons and cholesterol sulfate, may be close to values for these lipids in the plasma membranes of the fully differentiated cells of the stratum corneum. Such information is valuable for studies on the barrier function of the stratum corneum.

We thank Mrs. K. M. Boyce for technical assistance and Dr. Patricia Mann and Mr. R. H. Williams for the electron microscopy.

#### REFERENCES

- Gray GM, Yardley HJ: Different populations of pig epidermal cells: Isolation and lipid composition. *J Lipid Res* 16:441-447, 1975
- Hughes DE, Wimpenny JWT, Lloyd D: The disintegration of micro-organisms, *Methods in Microbiology*, vol 5B. Edited by JR Norris, DW Ribbons. New York and London, Academic Press, 1971, pp 1-54
- Wright BM, Edwards AJ, Jones VE: Use of a cell rupturing pump for the preparation of thymocyte subcellular fractions. *J Immunol Methods*, pp 281-296, 1974
- Gray GM, Tabiowo A, Trotter MD: Studies on the soluble and membrane bound amino acid 2-naphthylamidases in pig and human epidermis. *Biochem J* 161:667-675, 1977
- Lauter CJ, Solyom A, Trams EG: Comparative studies on enzyme markers of liver plasma membranes. *Biochim Biophys Acta* 266:511-523, 1972
- Taylor DG, Williams VM, Crawford N: Bis-(4-methylumbelliferyl) phosphate as a substrate for the surface membrane associated phosphodiesterase activity of pig platelets. *Biochim Biophys Acta* 465:667-671, 1977
- Strittmatter P: NADH-cytochrome b<sub>5</sub> reductase, *Methods in Enzymology* vol X. Edited by SP Colowick, NO Kaplan. New York, Academic Press, 1967, pp 561-565
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ: Protein measurement with the Folin phenol reagent. *J Biol Chem* 193:265-275, 1951
- Matoltsy AG, Matoltsy MN: The membrane protein of horny cells. *J Invest Dermatol* 46:127-129, 1966
- Block RJ, Weiss KW: *Amino Acid Handbook*. Springfield, Ill., C C Thomas, 1956, pp 278 and 341
- Gray GM, Yardley HJ: Lipid compositions of cells isolated from pig, human and rat epidermis. *J Lipid Res* 16:434-440, 1975
- Gray GM: Phosphatidyl-(N-acyl)-ethanolamine. A lipid component of mammalian epidermis. *Biochim Biophys Acta* 431:1-8, 1976
- Ugel AR, Chrambach A, Rodbard D: Fractionation and characterization of an oligomeric series of bovine keratohyalin by polyacrylamide gel electrophoresis. *Anal Biochem* 43:410-426, 1971
- Fairbanks G, Steck TL, Wallach DFH: Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochemistry* 10:2606-2617, 1971
- Zacharius RM, Zell TE, Morrison JH, Woodlock JJ: Glycoprotein staining following electrophoresis on acrylamide gels. *Anal Biochem* 30:148-152, 1969
- Gray GM, Yardley HJ: Mitochondria and nuclei of pig and human epidermis: Isolation and lipid composition. *J Invest Dermatol* 64:423-430, 1975
- Gray GM, White RJ, Majer JR: 1-(3'-O-acyl)- $\beta$ -glucosyl-N-dihydroxypentatriacontadienoylsphingosine, a major component of the glucosylceramides of pig and human epidermis. *Biochim Biophys Acta*, 528:127-137, 1978
- Baden HP, Lee LD, Kubilus J: The fibrous proteins of stratum corneum. *J Invest Dermatol* 67:573-576, 1976
- Skerrow CJ, Matoltsy AG: Chemical characterization of isolated epidermal desmosomes. *J Cell Biol* 63:524-530, 1974
- Rice RH, Green H: The cornified envelope of terminally differentiated human epidermal keratinocytes consists of cross-linked protein. *Cell* 11:417-422, 1977
- Yardley HJ, Goldstein DJ: Changes in dry weight and projected area of human epidermal cells undergoing keratinization as determined by scanning interference microscopy. *Br J Dermatol* 95:621-626, 1976
- Leroux A, Torlinski L, Kaplan JC: Soluble and microsomal forms of NADH cytochrome b<sub>5</sub> reductase from human placenta. *Biochim Biophys Acta* 481:50-62, 1977

## Announcement

A postgraduate course and workshop in Dermatopharmacology sponsored by the Department of Dermatology, University of California School of Medicine, San Francisco, will be held on August 25-27, 1978 in San Francisco. (CME Category credit hours = 20). For information write to Extended Programs in Medical Education, University of California, 1308 Third Avenue, San Francisco, California 94143.